

Microwave Assisted Oxidation of *N,N*-Dimethylaniline by Ditertiarybutyl Chromate in Non-Aqueous Media

RICHA KUMARI¹, SNIGDHA LAL¹, ANITA KUMARI GUPTA²,
SANGITA KATRE³ and HARI OM PANDEY⁴

¹Departement of Chemistry, Ranchi University,
Ranchi- 834008, Jharkhand, India

²Department of Chemistry, S S Memorial College, Ranchi University,
Ranchi-834008, India

³Department of Chemistry, C J Patel Arts, Commerce, Science & Post-graduate College,
Tirora, R T M Nagpur University, Maharashtra-441911, India

⁴Department of Chemistry, Ranchi University, Ranchi- 834008, Jharkhand, India
rahangdalebs@yahoo.co.in

Received 2 November 2013 / Revised 25 November 2013/ Accepted 2 January 2014

Abstract: Microwave assisted organic synthesis has grown into a powerful tool in the hands of chemists. In the present paper we report the study of products of oxidation of *N,N*-dimethylaniline by ditertiary butyl chromate in few non aqueous solvents like acetonitrile, tetrahydrofuran and dioxane. The substrate solution and ditertiary butyl chromate was taken in appropriate ratio, mixed, stirred and irradiated in domestic microwave for the specified periods. The solid products of different colour and compositions so obtained were studied by elemental analysis, FTIR curves and thermogravimetric mass loss pattern.

Keywords: *N,N*-Dimethylaniline, Ditertiarybutylchromate, Dioxane, Tetrahydrofuran

Introduction

Microwave as a source of energy was first invented by Percey Spencer in 1947¹. Gedye *et al.*, has the credit to introduce microwave in organic syntheses in 1986². Microwave assisted organic synthesis (MAOS) and microwave induced organic reaction enhancement (MORE) have developed into powerful tool for organic syntheses since then. In comparison to conventional energy source like heating *etc.*, microwave irradiation leads to greater efficiency, better yield, enhanced specificity and less pollution with less energy consumption. Lidstrom *et al.*,³ published a detailed review of the organic reactions carried out by the application of microwave.

N,N-Dimethylaniline is a yellowish brown toxic liquid insoluble in water. It is precursor to many commercially important dyes such as malachite green and crystal violet. The oxidation of *N,N*-dimethylaniline by cupric chloride or by chloranil gives *N-p*-dimethyl

aminobenzyl-*N*-methylaniline, 4,4'-bisdimethylaminodiphenyl methane, crystal violet and methyl violet⁴. The anodic oxidation of *N,N*-dimethylaniline at Pt electrode in acidic buffer leads to *N,N,N',N'*-tetramethylbenzidine⁵. In addition to the electro-oxidative coupling as above, the main oxidation product of *N,N*-dimethylaniline with ring retention is C₆H₅N(CH₃)₂O whereas oxidation with ring degradation may lead to variety of products.

Keeping in view, the increasing use of microwave irradiation as a source of clean energy⁶, in conformity with the 12 principles of green chemistry⁷, the authors have undertaken the oxidation of *N,N*-dimethylaniline with ditertiary butyl chromate in microwave in non-aqueous solvents. The choice of ditertiary butyl chromate⁸ out of many chromium(VI) based oxidants like Collin's reagent⁹, pyridinium chlorochromate¹⁰, pyridinium florochromate¹¹, tetramethyl ammonium florochromate¹², quinolinium chlorochromate¹³ etc. is based on its docility accompanied with versatility towards organic substrates. Moreover, the preparation of ditertiary butyl chromate is less cumbersome in comparison to the preparation of other variants of chromium (VI). The oxidation products may serve as ligands to form adducts or complexes of chromium in different oxidation states^{14,15}. The study of these products may lead to valuable generalizations with regards to the mechanism and other aspects of the reaction.

Experimental

The chemicals used were all A.R. grade. Only freshly prepared solutions were used. The oxidant was prepared by dissolving crystals of CrO₃ in minimum quantity of tert. Butyl alcohol followed by filtration and decantation to get clear brown solution. Sample-601 was prepared by mixing a solution of 1 g (0.01 mol) of chromium trioxide in 10 mL of tert. butyl alcohol and a solution of 1.2 mL (0.01 mol) of *N,N*-dimethylaniline in 10 mL of acetonitrile as per the requirement for 1:1 oxidant:substrate ratio. The other samples R-602, R-603 were prepared by mixing solution of 0.01 mol of chromium trioxide in tert. butyl alcohol and solution of 0.02 and 0.03 mols of substrate required for 1:2 and 1:3 oxidant:substrate ratio in acetonitrile as solvent for the substrate. Similarly, samples R-604, R-605 and R-605 were prepared by the same procedure using tetrahydrofuran as solvent for substrate and samples R-606 and R-608 in solvent dioxane for the same. In all the cases, the solutions were mixed stirred and then irradiated in LG household microwave oven, MG-3937 C (microwave frequency 2450 MHz) for the period specified in Table 1. The solid products obtained, were washed several times until clear washing before collecting them as solid samples with symbols as above. The estimation of chromium in the products was done volumetrically using K₂S₂O₈ (in excess), 0.1 N K₂Cr₂O₇ solution and 0.1 N Mohr's salt solution. Elemental analysis of carbon, hydrogen and nitrogen was carried out by Elemental Analyser (Heraeus Vario EL III Carlo Erba 1108). FTIR spectra of the products were recorded on FTIR Spectrophotometer (Shimadzu 8201 PC). TGA was done on Thermal Analysis System (Perkin Elmer Diamond TG/DTA).

Results and Discussion

On the basis of observations as in Table 1 and Table 2, following generalization could be made:

- The reaction is much efficient in acetonitrile as a solvent than either in THF or dioxane. In acetonitrile it takes only 10 sec. as compared to 20 sec. in other cases.
- The products R-607 and R-608, as obtained in case of dioxane as solvent for substrate are almost same except that the number of water molecules is slightly less in case of higher oxidant : substrate ratio.

- The oxidation product in all the cases from R-601 to R-608 is *N,N*-dimethylaniline-*N*-oxide which gets associated with reduced chromium to form solid products.
- In samples R-601, R-602 and R-603 which are obtained in case of acetonitrile as solvent, chromium has gone to oxidation state as low as II whereas in other cases it is mostly in most stable III state.
- As we go from sample R-601 to R-603, the number of water molecules increases *i.e.* less number of water molecules get associated in the products when the ratio of oxidant is more. This is observed in case of THF and dioxane as solvent as well.
- The colour of the products in almost all the cases is grey which may be due to charge transfer in the mixed oxide of chromium.
- As per formulations arrived at, the reaction did not results in any coupling of the rings as reported earlier^{3,4}.
- Recrystallization of the products in suitable solvent and the study of the crystals formed may be the further course of investigation.

Table 1. Reactants

Product Code	Substrate	Solvent	Oxidant	Oxidant:Substrate Ratio	Irradiation Time
R-601	DMA	Acetonitrile	TBC	1:1	10 sec
R-602	''	''	''	1:2	10 ''
R-603	''	''	''	1:3	10 ''
R-604	''	THF	''	2:1	20 ''
R-605	''	''	''	1:1	20 ''
R-606	''	''	''	1:2	20 ''
R-607	''	Dioxane	''	2:1	20 ''
R-608	''	''	''	1:1	20 ''

Table 2. Products

Product Code	Colour	Solubility in Water	Empirical Formula	Formulation
R-601	Ash Grey	Insoluble	Cr ₅ C ₉ H ₂₇ O ₁₄ N	2Cr ₂ O ₃ .CrO.HCOOH. C ₆ H ₅ N(CH ₃) ₂ O.4H ₂ O.
R-602	Dark Grey	''	Cr ₅ C ₈ H ₃₁ O ₁₇ N	2Cr ₂ O ₃ .CrO.C ₆ H ₅ N(CH ₃) ₂ O. 9H ₂ O.
R-603	Dark Grey	''	Cr ₇ C ₁₀ H ₄₆ O ₂₆ N	3Cr ₂ O ₃ .CrO.C ₆ H ₅ N(CH ₃) ₂ O 2HCOOH.11H ₂ O.
R-604	Dark Grey	''	Cr ₆ C ₁₃ H ₃₉ O ₂₂ N	3Cr ₂ O ₃ . C ₆ H ₅ N(CH ₃) ₂ O. 5HCOOH.2H ₂ O.
R-605	Ash Grey	''	Cr ₄ C ₈ H ₂₃ O ₁₆ N	2Cr ₂ O ₃ .C ₆ H ₅ N(CH ₃) ₂ O.9H ₂ O.
R-606	Dark Grey	Sparingly Soluble	Cr ₃ C ₉ H ₂₃ O ₁₆ N	Cr ₂ O ₃ .CrO. C ₆ H ₅ N(CH ₃) ₂ O. HCOOH.9H ₂ O.
R-607	Ash Grey	''	Cr ₄ C ₉ H ₂₇ O ₁₃ N	2Cr ₂ O ₃ .C ₆ H ₅ N(CH ₃) ₂ O. HCOOH.4H ₂ O.
R-608	Ash Grey	Insoluble	Cr ₄ C ₈ H ₂₆ O ₁₄ N	2Cr ₂ O ₃ .C ₆ H ₅ N(CH ₃) ₂ O. HCOOH. 5H ₂ O.

Acknowledgement

The authors are thankful to CIF, BIT, Mesra, Ranchi for elemental analysis, FTIR curves and Research & Development centre, Tejpur University, Assam for DTA-TGA analysis of the samples.

References

1. Spencer, Percy L, Methods of treating foodstuff. US. Patents- 2495429, Jan-24, 1950.
2. Gedye R, Smith F, Westaway K, Ali H, Balidsera L, Lorraine B and Rousell J, *Tetrahedron Lett.*, 1986, **27(3)**, 279-282; DOI:10.1016/S0040-4039(00)83996-9
3. Lidstrom P, Tierney J, Wathey B and Westman J, *Tetrahedron Report No.* 2001, **589(57)**, 9225-9283.
4. Lindsay J R Norman R O C and Walker W M, *J Chem Soc B: Phys Org.*, 1968, **0**, 66-71.
5. Mizoguchi T and Adams R N, *J Am Chem Soc.*, 1961, **84(11)**, 2058-2061; DOI:10.1021/ja00870a009
6. Loupy A, Petit A, Hamelin J, Françoise T B, Jacquault P and Mathe D, *Synthesis*, 1998, **09**, 1213-1234; DOI:10.1055/s-1998-6083
7. "The 12 principles of Green Chemistry" United States Environmental Protection Agency, Retrieved-31, 2006.
8. Oppenauer R V and Oberrauch H, (Univ. Innsbruck, Austria), *Anaus Assoc Guin Argentina*, 1949, **37**, 246-262.
9. Collins J C, Hess W W and Frank F J, *Tetrahedron Lett.*, 1968, **9(30)**, 3363-3366; DOI:10.1016/S0040-4039(00)89494-0
10. Corey E J and Suggs J W, *Tetrahedron Lett.*, 1975, **16(31)**, 2647-3650; DOI:10.1016/S0040-4039(00)75204-X
11. Bhattacharjee M N, Choudhuri M K, Dasgupta H S, Roy N and Khathing D T, *Synthesis*, 1982, **1(7)**, 588-590; DOI:10.1055/s-1982-29872
12. Kassae M Z, Mahjoub A R and Ghammani S, *Tetrahedron Lett.*, 2003, **44(24)**, 4555-4557; DOI:10.1016/S0040-4039(03)00989-4
13. Srinivasan R, Ramesh C V, Madhulatha W and Balasubramaniam K, *Indian J Chem.*, 1996, **35B**, 480-484.
14. Katre S, Gupta A K and Pandey H O, *Asian J Expt Sci.*, 2012, **26(1)**, 1-4.
15. Gupta A K and Pandey H O, *Asian J Expt Sci.*, 2012, **26(2)**, 43-46.